

EXHIBIT 3

OAR Chapter 333, Division 200

OREGON HOSPITAL TRAUMA TEAM ACTIVATION CRITERIA

Systolic blood pressure <90 mmHg; or
Respiratory rate <10 or >29 breaths per minute (<20 in infant aged <1 year); or
Need for ventilatory support; or
Glasgow Coma Scale <9; or
All penetrating injuries to head, neck, torso, and extremities proximal to elbow or knee; or
Chest wall instability or deformity (e.g. flail chest); or
Two or more proximal long-bone fractures; or
Suspected spinal cord injury with motor sensory deficit; or
Transfers requiring blood transfusions; or
Emergency physician's discretion.

YES



**Activate
Full Trauma
Team**

Full Trauma Team

- General Surgeon
- Emergency Physician
- Emergency Nurse(s)
- Laboratory
- Radiology
- Respiratory Therapist

Response times from
patient arrival:

Level I and II – 15 min.
Level III and IV – 30 min

Glasgow Coma Scale of 9 - 13; or
Crushed, degloved, mangled, or pulseless extremity; or
Amputation proximal to wrist or ankle; or
Open or depressed skull fracture; or
Suspected pelvic fracture; or
Falls

- Adults: >20 feet (one story is equal to 10 feet); or
- Children: >10 feet or two or three times the height of the child; or

High-risk auto crash

- Intrusion, including roof: >12 inches occupant site; >18 inches any site; or
- Ejection (partial or complete) from automobile; or
- Death in same passenger compartment; or
- Vehicle telemetry data consistent with a high risk of injury; or

Auto vs. pedestrian/bicyclist thrown, run over, or with significant (>20 mph) impact; or
Motorcycle or ATV crash >20 mph; or
EMS provider or receiving hospital judgment

YES



**Activate
Modified
Trauma Team**

Modified Trauma Team

- Emergency Physician
- Emergency Nurse(s)
- Laboratory
- Radiology

CO-MORBID FACTORS

Older Adults

- Risk of injury/death increases after age 55 years; or
- SBP <110 might represent shock after age 65 years; or
- Low impact mechanisms (e.g. ground level falls) might result in severe injury; or

Children

- Should be triaged preferentially to pediatric capable trauma centers; or

Pregnancy >20 weeks; or

Burns

- Without other trauma mechanism: triage to burn facility; or
- With trauma mechanism: triage to trauma center; or

Anticoagulants and bleeding disorders

- Patients with head injury are at high risk for rapid deterioration; or

Time sensitive extremity injury